

## *Sapphicorhynchus*, a new early Givetian rhynchonellid (brachiopod) genus from western New York State, USA, and Sapphicorhynchidae, n. fam.

by Paul SARTENAER

SARTENAER, P., 2007 - *Sapphicorhynchus*, a new early Givetian rhynchonellid (brachiopod) genus from western New York State, USA, and Sapphicorhynchidae, n. fam. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, 77: 41-61, 1 pl., 2 text-figs, 1 table, Brussels, October 15, 2007 – ISSN 0374-6291.

### Abstract

A new genus, *Sapphicorhynchus*, type species *S. sappho* (HALL, 1860), is described from the early Givetian of western New York State. The presence of the species in older and younger strata of New York, in other States of the United States of America, in the Province of Ontario (Canada), and outside North America is rejected. The genus is designated the type for the new family Sapphicorhynchidae. It is compared to the emended middle Eifelian genus *Oligothycherhynchus* SARTENAER, 1970, type species *O. hexatomus* (SCHNUR 1851), and the early Givetian genus *Cupularostrum* SARTENAER, 1961, to which *Sapphicorhynchus sappho* is assigned in recent literature.

**Keywords:** Sapphicorhynchidae, *Sapphicorhynchus*, rhynchonellids, brachiopods, early Givetian, western New York.

### Résumé

L'auteur fonde un nouveau genre, *Sapphicorhynchus*, avec *S. sappho* (HALL, 1860) du Givetien Inférieur de la partie occidentale de l'État du New York comme espèce-type; la présence de l'espèce dans des couches plus anciennes et plus jeunes de cet État, dans d'autres États des États-Unis d'Amérique, dans la Province de l'Ontario au Canada et en dehors de l'Amérique du Nord est rejetée. Le genre, désigné comme type de la nouvelle famille Sapphicorhynchidae, est comparé à deux autres: *Oligothycherhynchus* SARTENAER, 1970, emend. de l'Eifélien moyen, avec *O. hexatomus* (SCHNUR 1851) comme espèce type et *Cupularostrum* SARTENAER, 1961, auquel *Sapphicorhynchus sappho* a été attribué dans la littérature récente.

**Mots-clefs:** Sapphicorhynchidae, *Sapphicorhynchus*, Rhynchonellides, Brachiopodes, Givetien Inférieur, New York occidental.

### Introduction

The revision of the taxonomical status and the stratigraphical range of key rhynchonellid taxa of the New York Devonian, that started by the author with the late Givetian *Tullypothyridina venustula* (HALL, 1867) in 2003 and the late Eifelian *Cherryvalleyrostrum limitare* (VANUXEM, 1842) in 2004, is carried on with the re-examination of the early Givetian *Rhynchonella sappho* HALL, 1860.

The extensive stratigraphical range - Upper Helderberg to Lower Carboniferous - of the species, which has successively been assigned to the genera *Rhynchonella* FISCHER DE WALDHEIM, 1809, *Stenosisma* CONRAD, 1839 (exceptionally), *Camarotoechia* HALL & CLARKE, 1893 (for about a century), and *Cupularostrum* SARTENAER, 1961, excludes the possibility to deal with the various forms to which its name has been given. It would lie outside the scope of the present study, which rests on the examination of the type species, field work, collections examined in various museums and scientific institutions, and scrutiny and pondering on the abundant literature on the subject.

### Systematic palaeontology

#### Sapphicorhynchidae n. fam.

*Type genus:* *Sapphicorhynchus* n. gen.

#### Diagnosis

Shell of medium to medium-large size with strongly dorsibiconvex profile and subcircular to subpentagonal outline; top of shell posterior to front; short ventral interarea, well marked lunulae; angle of the cardinal commissure wide. Sulcus wide, moderately deep, fold



low, start at a short distance from beaks; costae well marked, simple, regular, in moderate number, start at the beaks, and strongly indent the commissure; parietal costae sometimes present. Delicate internal structures. Undivided hinge plate, crura raduliform. Dental plates separated from the wall by wide umbonal cavities; septum long; septalium wide and deep, amphora-shaped; connectivum covering anterior part of septalium.

#### Remarks

The diagnostic combination of characters allows separation of the family from all known families, and in particular from the family Trigonirhynchiidae SCHMIDT, 1965 here discussed.

Various reasons explain the remarkable prosperity of the family Trigonirhynchiidae. The main one, which has already been pointed out by SARTENAER (2001, p. 208), is the inclusion in the short diagnosis of the family of many alternative characters (see Fig. 2); the above-mentioned diagnosis does not contain any such characters. Consequently: 1) the family becomes an "accommodating drawer" and the few non-alternative remaining characters are shared with other families; 2) some other characters are omitted; 3) undue emphasis is placed on one or two or few characters; 4) genera are easily transferred from one family to another. These four points are exemplified.

Nine genera were added by MCLAREN in SCHMIDT & MCLAREN (1965, pp. H559-562), three of them questionably (*Lepidocycloides* NIKIFOROVA in NIKIFOROVA & ANDREEVA, 1961, *Nekhoroshevia* BUBLICHENKO, 1956, and *Rostricellula* ULRICH & COOPER, 1942) to the four genera (*Cupularostrum* SARTENAER, 1961, *Nymphorhynchia* RZHONSNITSKAYA, 1956, *Ptychomaletoechia* SARTENAER, 1961, and *Trigonirhynchia* COOPER, 1942) originally assigned to the family Trigonirhynchiidae by SCHMIDT (1965, pp. 2, 3). Thirty years later the family numbered some 50 genera. In 2007 the number rose to about 80, making it the "richest" rhynchonellid family, the second "richest" being the Leiorhynchidae STAINBROOK, 1945. Out of them 54 (53 in 1996, 54 in 2002), 21 for the first time, were listed by SAVAGE (1996, table 3, p. 256; 2002, pp. 1052-1077) and distributed among five subfamilies: Hemitoechiinae SAVAGE, 1996 (17 genera), Ripidiorhynchinae SAVAGE, 1996 (four genera), Rostricellulinae ROZMAN, 1969 (four genera in 1996, five in 2002), Trigonirhynchiinae SCHMIDT, 1965 (24 genera), and Virginiatiinae AMSDEN, 1974 (four genera).

It is not appropriate to single out a few characters or to decree that some are major and others minor. Characteristics of a family or a genus must be considered as a whole. Why should features such as the following be overlooked, omitted or rejected: location of top of shell; apical angle; depth, width and point of origin of sulcus and fold; regularity, strength, point of origin, number of costae; presence or absence of parietal costae; shape, depth, strength of the septalium; and the serration (undulation, indentation) of the commissure. In the case of *Trigonirhynchia fallaciosa* (BAYLE, 1878), the type species of *Trigonirhynchia*, its evident "globular, subcuboidal" shape mentioned by OEHLERT (1884, p. 420) in the first description of the species is not included in the diagnosis of the genus, and neither is the great deepness of the septalium illustrated by SCHMIDT (1965, fig. 1, p. 4) and reproduced by SCHMIDT in SCHMIDT & MCLAREN (1965, fig. 428, 1f-q, p. H559) and SAVAGE (2002, fig. 710, 1e-n, p. 1053). Furthermore, the internal structure of *T. fallaciosa* cannot be considered as fully known; in order to better understand the transverse serial sections given by SCHMIDT (1965), the author made sections in another topotype that show a stout septalium and connectivum, and dental plates separated from the wall by narrow umbonal cavities.

When few characters (sometimes one or two) are considered, the classification may move in various directions depending upon the weight put on them. Thus, AMSDEN (1974, p. 68) divided the family Trigonirhynchiidae into two subfamilies: Trigonirhynchiinae (forms with a fold and sulcus), and Virginiatiinae AMSDEN, 1974 (forms lacking a fold and sulcus). ROZMAN (1969, pp. 86, 94-96) erected the family Rostricellulinae embracing three genera (*Evenkorhynchia* ROZMAN, 1969, *Lepidocycloides*, and *Rostricellula* ULRICH & COOPER, 1942) to which SAVAGE (1996, table 3, p. 256; 2002, p. 1065) added the genera *Azamella* LAURIE, 1991 in 1996 and *Plectothyrella* TEMPLE, 1965 in 2002. JIN *et al.* (1993, p. 54) adopted the subfamilies Trigonirhynchiinae and Rostricellulinae as the two divisions of the family Trigonirhynchiidae, and assigned eight genera to the former and 16 genera to the latter, but considered as distinctive the "septalial cavity covered completely or anteriorly by medially coalescent plates" (Trigonirhynchiinae), and the "septalial cavity commonly restricted, but not completely covered by medially projected crural ridges" (Rostricellulinae). BARANOV (1989, p. 42) had already proposed to follow a similar line, but with a different approach;



assuming that the structure of the cardinalia of the family Trigonirhynchiidae “did not undergo important changes in the course of evolution since the Ordovician”, he stated that divided and undivided hinge plates “interpreted by scientists as a taxonomical character of generic and family level” appeared independently in different phylogenetic branches, and that probably the “separation of Ancillotoechia-like [i.e. covered septalium] genera took place a few times in the course of iterative evolution”. Meanwhile, HAVLÍČEK & ŠTORCH (1990, p. 125) considered that the subfamily Rostricellulinae “was very close to the Oligorhynchiidae [COOPER, 1956]” “judging from the inner and outer morphology”, and it is to this family that GARCÍA-ALCALDE (1998, p. 769) assigned five genera included by SAVAGE (1996) in the subfamily Trigonirhynchiinae [*Ancillotoechia* HAVLÍČEK, 1959, *Aratoechia* HAVLÍČEK, 1982, *Myrmirhynx* HAVLÍČEK, 1982, *Sufetirhynchia* HAVLÍČEK, 1982, and *Iberirhynchia* DROT & WESTBROEK, 1966; *Ancillotoechia*, *Myrmirhynx* and *Sufetirhynchia* were already assigned to the subfamily by JIN *et al.* (1993, p. 54)] plus a new genus *Tectogonotoechia* GARCÍA-ALCALDE, 1998 not mentioned by SAVAGE (1996, 2002).

Moving genera in and out of the family Trigonirhynchiidae is a common practice that is bound to last in the next future, e.g. *Werneckeella*, LENZ 1971 put by BRICE in BRICE *et al.* (2000, fig. 1, p. 68) in the family Trigonirhynchiidae is now included in the subfamily Leiorhynchinae STAINBROOK, 1945. Such shifts emphasize the deficiency and precariousness of the present classification. They can lead to the shrinking of a family to a few genera, when not only to its type genus. This has been the case of the family Camarotoechiidae SCHUCHERT & LE VENE, 1929, in which SCHMIDT & McLAREN (1965, pp. H579-584) included twenty genera distributed among two subfamilies: Camarotoechiinae SCHUCHERT & LE VENE, 1929 (16 genera) and Septalariinae HAVLÍČEK, 1960 (four genera).

To conclude, many discrepancies and/or contradictions result from the misleading diagnosis of the family Trigonirhynchiidae. Only one example is given here. What has the tiny lower-middle Eifelian genus *Tetratomia* SCHMIDT, 1941 in common with the medium-large to large Pragian [not Emsian as stated by SAVAGE (1996, table 3, p. 256; 2002, p. 1070)] genus *Losvia* BREIVEL & BREIVEL, 1976, the former assigned by SAVAGE (2002, p. 1070) to the subfamily Trigonirhynchiinae, the latter to the subfamily Hemitoechiinae SAVAGE, 1996? (*Tetratomia* has a

narrow and well marked sulcus and fold, an undivided hinge plate, no septalium, etc. *Losvia* has a weak sulcus and fold, a very low (almost none) tongue, an uncovered septalium, etc.)

### *Sapphicorhynchus* n. gen.

#### *Derivatio nominis*

Σαπφικός, ἡ, όν (Greek, adjective) = sapphic. The name draws attention to the name given to the species, now type species, by HALL (1860).

*Type species: Rhynchonella sappho* HALL, 1860.

It is not to be excluded that some forms from the lower and middle Ludlowville Formation (Middle Givetian) of western and central New York State (Centerfield Limestone, Ledyard Shale, Wanakah Shale, and Otisco Shale Members) that have been assigned to the species could belong to the genus, if not to the species. Such forms have neither been figured nor described, and their examination would require more material than the one at hand.

#### *Diagnostic features*

Shell of medium to medium-large size, outline subcircular to subpentagonal, greatest thickness posterior to front. Ventral interarea short, angle of the cardinal commissure wide. Sulcus and fold start at a short distance from the beaks, sulcus moderately deep, fold low, forming a moderately high transverse-trapezoidal tongue, top of tongue lower than the maximum shell thickness. Costae in moderate number, start at the beaks. Parietal costae sometimes present. Internal structures delicate. Dental plates separated from the wall by wide umbonal cavities. Hinge plate undivided, connectivum covering anterior part of septalium, septum long, septalium wide and deep, amphora-shaped, elongated raduliform crura.

#### *Description*

Shell medium- to medium-large-size, profile strongly dorsibiconvex (dorsal valve considerably thicker than ventral valve, the flanks of which are flattening near the front in medium-large specimens); ventral valve thickest at 1/4 to 1/3 shell length, dorsal valve thickest anterior to mid-length. Shell outline subcircular to subpentagonal (length about 3/4 to 4/5 width); half barrel-shaped in frontal view. Postero-lateral and lateral commissures sharp, strongly indented by the costae. Anterior commissure fused into the wall formed by the extremities of the costae. Wide angle of the cardinal commissure. Top of shell posterior



to front. Elongated concave lunulae present in both valves with zero to two costae (two in the lectotype) on each of them; they are defined by faint beak ridges near the beaks. Ventral beak acute, slightly to strongly incurved; umbo low. Ventral interarea short; deltidial plates narrow. Sulcus and fold well marked, start imperceptibly at a short distance from the beaks. Sulcus wide, moderately deep and well delineated towards margin; bottom flat to slightly convex, extended dorsally as a moderately high tongue with subtrapezoidal outline, recurving slightly posteriorly in its uppermost part. Fold remains low throughout; top gently convex. In transverse profile, dorsal valve high, outer flanks deflected ventrally to be almost vertical near commissure. Median and lateral costae, in moderate number, start from the beaks [Remark: number of costae is not related to the ontogeny as implied by the founder of the species (HALL, 1860, p. 87; 1867, p. 340)]. Costae well marked, moderately high, wide, rounded, simple, and regular. Costae (with the exception of the postero-lateral ones) interlock in high zigzags at the commissure. Parietal costae, when present, reach and indent the commissure.

Internal structures delicate. Dental plates subparallel posteriorly, becoming convergent anteriorly; umbonal cavities wide; teeth short, with wrinkled dorsal face. Inner socket ridges low. Septum long; septalium deep and wide (deeper than wide), amphora-shaped; hinge plate undivided, outer hinge plates flat to slightly convex passing without sign of crural bases into long crura; crura raduliform, Phrygian cap-shaped in section in their distal part where they are slightly curved ventrally; connectivum covering anterior part of septalium, and extending beyond.

#### Growth

With increasing size specimens become wider (transverse), and their angle of the cardinal commissure opens. Ventral lateral costae are often fan-wise in medium-large specimens; the term "flabelliform" used by HALL (1860, p. 87; 1867, p. 340) applies to such specimens, which HALL (1867, explanation of pl. 54, figs 38-42) and HALL & CLARKE (1893, explanation of pl. LVII, figs 10-14) considered as "large and characteristic" or "large and typical" (see Pl. 1, Figs 1-5).

#### Comparisons

The Lower Givetian (Upper Pompey Member of the Skaneateles Formation) genus *Cupularostrum* SARTENAER, 1961 has few characters in common

with *Sapphicorhynchus* n. gen. It is only mentioned here because *S. sappho* has been included in it by GRIESEMER (1965, pp. 268-270), and since then, by North American and non-American authors (see Synonymy) [Remark: Three species were already assigned to the genus in 1964: the late Silurian *Cupularostrum litchfieldense* (SCHUCHERT, 1903) and the Lower Devonian *C.? semiplicata* (CONRAD, 1841) by BERDAN (1964, p. 16), and a new species, *C. cantabricum*, collected by WESTBROEK (1964, pp. 243, 244-246) near the Frasnian/Famennian boundary]. JOHNSON & PERRY (1976, pp. 622-624), who discussed the genus at length, ignored GRIESEMER's paper in which, with the exception of the type species, *C. recticostatum*, SARTENAER, 1961, none of the twelve species and variety attributed to *Cupularostrum* belong to that genus [*Camarotoechia ambigua*, FAGERSTROM 1961, *C. cedarensis*, STAINBROOK, 1942, *C. congregata* var. *parkheadensis* CLARKE & SWARTZ, 1913, *Rhynchonella depressa* KINDLE, 1901, *Atrypa eximia*, HALL, 1843, *Camarotoechia formosensis* FAGERSTROM, 1961, *C. gregeri* BRANSON, 1923, *Rhynchonella orbicularis* HALL, 1860, *R. (Stenocisma) cf. prolifica* HALL, 1867, *Rhynchonella sappho*, *Camarotoechia scitulus* CLELAND, 1911, and *Rhynchonella tethys* BILLINGS, 1860]. In general terms, *Cupularostrum*, to which some 50 species and subspecies + some 40 forms in open nomenclature have been assigned at one time or another, almost all of them incorrectly, is poorly understood and its alleged cosmopolitan status advocated by JOHNSON & BOUCOT (1973, p. 95), NORRIS (1979, p. A246), BRICE (in RACHEBOEUF *et al.*, 2001, p. 147), and SAVAGE (2002, p. 1056) is disputable. So is also its notable - Upper Silurian to Lower Viséan - and variable stratigraphical range, e.g. Upper Emsian to Middle Frasnian (SAVAGE, 1996, table 3, p. 256; 2002, p. 1056), latest Eifelian to earliest Frasnian (BRICE in BRICE *et al.*, 2000, fig. 1, p. 68). SAVAGE *et al.* (1979), although they incorrectly accepted the presence of "*Cupularostrum*" in the Lochkovian, Pragian, and Emsian (table, p. 175), were right when they wrote (p. 195) that "*Cupularostrum*" "is a poorly known Lower Devonian genus and the occurrences listed [table, p. 175] may include more than one genus". MERGL *et al.* (2001, p. 178) were also certainly entitled to call *Cupularostrum* a "basket" genus.

*Cupularostrum* can easily be separated from *Sapphicorhynchus* n. gen. by its subglobular aspect and its thick and robust internal features (dental plates, septum, hinge plate, connectivum). Other



distinctive characters are a smaller size (the largest adult specimen of *C. recticostatum* reaches the size of the smallest adult specimen of *Sapphicorhynchus sappho*); a shell outline always subcircular or oval; a relatively higher ventral valve; the maximum thickness of the shell located more posteriorly; different l/w ratios (0.81 to 1 against 0.71 to 0.88 for *S. sappho*); narrower sulcus and fold; the uppermost part of tongue not slightly recurved posteriorly, but tending sometimes to become vertical; a different general costal formula ( $\frac{4 \text{ to } 5}{3 \text{ to } 4}$ ; 0 to  $\frac{1-1}{1-1}$ ;  $\frac{8 \text{ to } 10}{9 \text{ to } 11}$  against  $\frac{5 \text{ to } 6}{4 \text{ to } 5}$ ; 0 to  $\frac{1-1}{1-1}$ ;  $\frac{6 \text{ to } 8}{7 \text{ to } 9}$  for *S. sappho*); generally narrower costae; parietal costae rarely present; a smaller angle of the cardinal commissure (97° to 115° against 115° to 128° for *S. sappho*); and lower inner socket ridges.

Before comparing *Sapphicorhynchus* n. gen. with *Oligoptycherhynchus* SARTENAER, 1970, the following remarks have to be made. When establishing the genus *Oligoptycherhynchus*, with type species *O. hexatomus* (SCHNUR, 1851) from the middle Eifelian (Ahrdorf Schichten) of the Gerolsteiner Mulde, SARTENAER (1970, pp. 20-21) also included: *Terebratula daleidensis* ROEMER, 1844 from the upper Emsian (Wiltz Schichten) of the Daleiden "Muldengruppe", *T. elliptica* SCHNUR, 1853 from the middle Eifelian (Gondelsheimer Schichten) of the Gerolsteiner Mulde, *Rhynchonella imitatrix* FUCHS in SPIESTERSBACH & FUCHS, 1909 from the upper Emsian (Remscheider Schichten) of the Bergisches Land, and *Camarotoechia hexatoma wetteldorfensis* SCHMIDT, 1941 from the uppermost Emsian (Heisdorf Schichten) of the Prümer Mulde. Field work carried out since then in the Eifel area and in the Bergisches Land, and the study of collections housed in various museums and scientific institutions have convinced the author that the species and subspecies had a considerably shorter stratigraphical range (see above), and that they had to be rejected from the genus. Consequently, the wide range (middle Siegenian to Eifelian) advocated by the author in 1970 for *Oligoptycherhynchus* has become unacceptable, and the genus, now defined by its type species only, has to be emended. The emended diagnosis is as follows (changes in the initial diagnosis are in italics):

*Medium size. outline subcordiform. Uniplicate. Inequivalve. Top of the dorsal valve at front, Commissure sharp. Well marked sulcus and fold, starting at a short distance from the beaks. Sulcus moderately deep with flat to slightly convex bottom,*

*wide at front. High fold with slightly convex top. High subrectangular tongue with sharp borders, sometimes recurved posteriorly in its upper part. Ventral beak slightly to strongly incurved. Narrow ventral interarea. Costae in moderate number, regular, simple, moderately elevated to elevated, angular or angular with rounded top, starting at the beaks. Parietal costae common, apical angle wide. Moderately thick shell. Moderately thick dental plates, diverging in the apical region, becoming parallel to convergent anteriorly. Long and moderately thick septum. Deep septalium, amphora-shaped in transverse serial sections, covered by a moderately robust connectivum. Hinge plate composed of two horizontal to slightly concave parts. Inner ridges of dental sockets high. Short and robust teeth.*

Another point that has commonly been overlooked is that the specimen of *Oligoptycherhynchus hexatomus* designated by SCHMIDT (1941, p. 9) as the representative of the species is not a specimen chosen in the original SCHNUR's (1851, 1853) collection. It also comes neither from the *locus typicus*, nor from the *stratum typicum*, which are the "Kalk zu Pelm und Gees", i.e. the Gees Horizon of the middle Eifelian Ahrdorf Schichten of the Gerolsteiner Mulde in the Eifel area. Assuming that the two specimens figured by SCHNUR (1853, pl. 23, figs 2a-g) were lost, SCHMIDT (1941, p. 9) took the unfortunate decision to consider a specimen coming from N. of Üxheim ("n. Ü. von Üxheim") as the representative of the species and considered by her as approaching "am besten der Abbildung bei SCHNUR" the specimen (figs 2a-e) figured by SCHNUR. But Üxheim is located in a different syncline, the Hillesheimer Mulde, than Pelm and Gees. Moreover, SCHMIDT's (1941, p. 9) statement that the "Horizont des Üxheimer Fundorts ist ja auch bei Gees vertreten" is contradicted by WEDEKIND (1924, pp. 86-90) and STRUVE in HOTZ *et al.* (1955, p. 59), who observed that the alleged Geeser Horizont N. of Üxheim is the Kirberg Member of the Lower Nohner Schichten (lower Eifelian).

The middle Eifelian genus *Oligoptycherhynchus* and *Sapphicorhynchus* n. gen. have many characters in common, e.g. a strongly biconvex profile; sharp postero-lateral and lateral commissures, strongly indented by the costae; an anterior commissure fused into the wall formed by the extremities of the costae; a short ventral interarea; well marked sulcus and fold, starting imperceptibly at some distance from the beaks; a wide, moderately deep sulcus, with flat to slightly convex bottom; a tongue slightly recurved posteriorly in its uppermost part; the top of fold gently convex; a



moderate number of well marked, moderately high, wide, rounded, simple, regular costae, starting at the beaks; dental plates converging anteriorly; wide umbonal cavities; a long septum; a deep amphora-shaped septalium; a connectivum covering the anterior part of the septalium and extending beyond (undivided hinge plate); short robust teeth; and long raduliform crura.

Many characters make *Sapphicorhynchus* n. gen. distinct from *Oligoptycherhynchus*: a smaller thickness; a greater width (this is only valid for middle-large specimens of *Sapphicorhynchus sappho*); the maximum thickness located posterior to front; better marked lunulae; a moderately high tongue with transverse-subtrapezoidal outline; a moderately high fold; ventral lateral costae arranged fan-wise (this is only valid for middle-large specimens of *S. sappho*); parietal costae less present, reaching and indenting the commissure; a wider angle of the cardinal commissure; a thinner shell; more delicate internal structures; horizontal or slightly convex outer hinge plates passing without sign of crural bases to Phrygian cap-shaped crura in section made at their distal end; and lower inner socket ridges. Finally, some specimen of *S. sappho* may reach larger sizes than the largest specimens of *Oligoptycherhynchus hexatomus*.

### *Sapphicorhynchus sappho* (HALL, 1860)

Plate 1, Figs 1-33; Text-figure 1

- 1860 – *Rhynchonella sappho* (n.s.) – HALL, p. 87.  
 1867 – *Rhynchonella* (*Stenocisma*) *sappho* – HALL, pp. 340–341, pl. 54, figs 34–43.  
 non 1867 – *Rhynchonella* (*Stenocisma*) *sappho* – HALL, pl. 54, fig. 33.  
 non 1867 – *Rhynchonella* (*Stenocisma*) *sappho* var. – HALL, p. 354, pl. 55, figs 47–52.  
 non 1884 – *Rhynchonella Sappho* – WILLIAMS, p. 23.  
 non 1887 – *Rhynchonella Sappho* – WILLIAMS, table, p. 27.  
 non 1887 – *Rhynchonella?* *Sappho*, var. – WILLIAMS, p. 62.  
 non 1887 – *Rhynchonella sappho*, var. – WILLIAMS, pp. 74, 77.  
 non 1887 – *Rhynchonella Sappho*, var. – WILLIAMS, p. 88.  
 non 1887 – varieties of *R. Sappho* – WILLIAMS, p. 88.  
 non 1888 – *Rhynchonella sappho*, Hall – HERRICK, pp. 24, 25, 26, 40, table, p. 98, pl. VII, fig. 25.  
 non 1888 – *Rhynchonella sappho* (varieties of...) – HERRICK, pp. 39, 40.  
 non 1888 – *Rhynchonella sappho* (conditions of...) – HERRICK, p. 40.  
 non 1888 – *Rhynchonella sageriana* (= var. of *R. sappho* ?) – HERRICK, pl. II, fig. 15.  
 non 1888 – *Rhynchonella sappho* (Cf. *cooperi*) – HERRICK, pl. V, fig. 1.  
 non 1889 – *Rhynchonella sappho* – LESLEY, p. 900.  
 1890 – *Stenoschisma sappho*. (*Rhynchonella sappho*) Hall – LESLEY, p. 1060, figs 34–43, p. 1060 (= pl. 54, figs 34–43 in HALL, 1867).  
 non 1890 – *Stenoschisma sappho*. (*Rhynchonella sappho*) Hall – LESLEY, fig. 33 = pl. 54, fig. 33 in HALL, 1867.  
 non 1890 – *Stenoschisma sappho*. (*Rhynchonella sappho*) Hall, a variety ? – LESLEY, p. 1060.  
 non 1890 – *Stenoschisma sappho*. (*Rhynchonella sappho*) Hall – LESLEY, p. 1060.  
 1893 – *Rhynchonella Sappho*, Hall – HALL & CLARKE, pl. LVII, figs 10–14 (= pl. 54, figs 38–42 in HALL, 1867).  
 e.p. 1893 – *Rhynchonella Sappho*, Hall – HALL & CLARKE, p. 192.  
 non 1893 – *Rhynchonella Sappho* – HERRICK in HALL & CLARKE, p. 192.  
 non 1896 – *Rhynchonella sappho* – WILLIAMS in KINDLE, p. 47.  
 ? 1897 – *Rhynchonella* (*Camarotoechia*) *sappho*, HALL – PROSSER, p. 98.  
 non 1898 – *Camarotoechia Sappho*, Hall – WHITEAVES, pp. 386, 387.  
 non 1898 – *Camarotoechia Sappho* (Hall) – WHITEAVES, p. 416.  
 non 1899 – *Camarotoechia sappho* Hall (?) – GIRTY, table, p. 485, pp. 541–542.  
 non 1899 – *Camarotoechia sappho* (?) – GIRTY, pp. 490, 492.  
 e.p. 1899 – *Camarotoechia sappho* – GIRTY, p. 492.  
 1899 – *Camarotoechia sappho* Hall – GRABAU, fig. 140, p. 230 (= pl. 54, figs 38–42 in HALL, 1867).  
 non 1899 – *Camarotoechia sappho*. Hall. – GRABAU, p. 230.  
 non 1899 – *Camarotoechia sappho* Hall – KINDLE, pp. 13, 15, 16, 19, 28, 31, 58.  
 non 1899a – *Camarotoechia Sappho*, Hall – GRABAU, pp. 282, 296, 297, 299, 304, 308, 314, 318, table.  
 non 1899b – *Camarotoechia Sappho*, Hall – GRABAU, pp. 282, 296, 297, 299, 304, 308, 314, 318, table.  
 1901 – *Camarotoechia sappho* Hall – CLARKE, pp. 132, 134.  
 1901 – *Camarotoechia sappho* Hall – WOOD, pp. 150, 151, 162, 175, 179.  
 1901 – *Camarotoechia sappho* (Hall) – WOOD, p. 162.  
 non 1901 – *Camarotoechia sappho* Hall – KINDLE, pp. 568, 584, pl. VII, figs 4, 4a.  
 non 1902 – *Camarotoechia sappho* (Hall) – SHIMER & GRABAU, table, p. 182.  
 non 1903 – *Camarotoechia sappho* ? – BUTTS, p. 993.  
 non 1903 – *Camarotoechia sappho* Hall – BUTTS, pp. 994–995.  
 non 1903 – *Camarotoechia sappho* Hall – CLELAND, p. 42, appendix, p. 97.  
 non 1904 – *Camarotoechia sappho* Hall – RAYMOND, pp. 82, 151, 162.  
 non 1905 – *Camarotoechia sappho* Hall – CLARKE, p. 64.  
 non 1905 – *Rhynchonella* (*Camarotoechia*) cf. *Sappho* HALL – THOMAS, table, p. 286.  
 non 1905 – *Camarotoechia sappho* – KINDLE in WILLIAMS & KINDLE, pp. 21, 25, 32, 35, 120, table, p. 122.  
 non 1905 – *Camarotoechia sappho* – WILLIAMS in WILLIAMS & KINDLE, chart between p. 54 and p. 55.  
 non 1908 – *Rhynchonella Sappho* Hall – KNOD, p. 552.  
 non 1908 – *Camarotoechia* aff. *sappho* – GIRTY in KINDLE, p. 92.  
 non 1908 – *Camarotoechia sappho* – GIRTY in KINDLE, pp. 92, 98.  
 non 1909 – *Camarotoechia sappho* Hall? – STAUFFER, p. 122.  
 non 1909 – *Camarotoechia sappho* Hall – STAUFFER, table, p. 163.  
 1909 – *Camarotoechia sappho* Hall – GRABAU & SHIMER, figs 354 c,d, p. 287 = pl. 54, figs 39, 40 in HALL, 1867.  
 e.p. 1909 – *Camarotoechia sappho* Hall – GRABAU & SHIMER, p. 288.  
 1911 – *Camarotoechia sappho* – CLELAND, p. 25.  
 non 1912 – *Camarotoechia sappho* Hall var. – PROSSER, p. 90.



- non 1912 – *Camarotoechia sappho* – PROSSER, p. 91.  
 non 1912 – *Rhynchonella sappho* – PROSSER, p. 118.  
 e.p. 1913 – *Rhynchonella Sappho* Hall – LA TOUCHE, p. 246.  
 non 1913 – *Camarotoechia sappho* Hall – PROSSER in PROSSER et al., p. 77.  
 non 1913 – *Camarotoechia sappho* Hall – PROSSER in PROSSER & KINDLE, pp. 174–175, pl. XV, fig. 4.  
 e.p. 1913 – *Camarotoechia sappho* Hall – PROSSER in PROSSER et al., table, p. 106.  
 non 1915 – *Camarotoechia sappho* Hall – STAUFFER, pp. 139, 144, 159, 171, 178, 233.  
 non 1922 – *Camarotoechia sappho* Hall – BRANSON & WILLIAMS, p. 131, pl. 34, figs 15–17.  
 non 1922 – *Camarotoechia sappho* (Hall) – BRANSON & WILLIAMS, pp. 143–144.  
 non 1929 – *Camarotoechia* cf. *sappho* – SAVAGE, pp. 112, 249.  
 e.p. 1929 – *Camarotoechia sappho* – COOPER, pp. 73, 75, 81, 161, 183, 208, 210, 228, 232, 251.  
 non 1930 – *Camarotoechia sappho* (Hall) – CASTER, pp. 152, 155, 156, 157–161, pl. 33, figs 3, 14–18, pl. 34, figs 7–17 (synonymia pro parte).  
 ? 1930 – *Camarotoechia sappho* (some variety of...) – WILLIAMS in CASTER, p. 156.  
 e.p. 1930 – *Rhynchonella sappho* – HALL in CASTER, p. 158.  
 non 1930 – *Camarotoechia sappho* (minor variations of...) – CASTER, p. 159.  
 non 1930 – *Camarotoechia sappho* – CASTER, pp. 164, 174, 180.  
 1930 – *Camarotoechia sappho* (Hall) – CASTER, pl. 33, figs 7, 12, 13.  
 non 1930 – *Camarotoechia sappho* (Hall?) – CASTER, pl. 33, figs 8–10.  
 non 1930 – *Camarotoechia sappho* ? – CASTER, pl. 35, fig. 22.  
 non 1930 – *Rhynchonella sappho* Hall – FERUGLIO, table, p. 81, p. 85.  
 e.p. 1931 – *Rhynchonella (Camarotoechia) sappho* – GRABAU, p. 106.  
 non 1931 – *Camarotoechia sappho* – LAUDON, p. 253.  
 1931 – *Camarotoechia sappho* – GOLDRING, p. 391.  
 non 1932 – *Camarotoechia sappho* – VON ENGELN, p. 58.  
 non 1933 – *Camarotoechia* cf. *sappho* Hall – NALIVKIN, pp. 203, 204, 210–211.  
 non 1934 – *Camarotoechia sappho* Hall – CASTER, p. 161.  
 non 1935 – *Camarotoechia sappho* (Hall) – COOPER & WILLIAMS, pp. 790, 791, 813, list, p. 856.  
 non 1935 – *Camarotoechia* cf. *sappho* – SMITH, p. 104.  
 non 1935 – *Camarotoechia sappho* ? – SMITH, pp. 106, 109.  
 non 1935 – *Camarotoechia sappho* (Hall) – SMITH, pp. 110, 111.  
 non 1935 – *Camarotoechia sappho* – CHADWICK, pp. 317, 325, 331, 336.  
 non 1935 – *Camarotoechia sappho* (?) – CHADWICK, p. 331.  
 non 1937 – *Camarotoechia sappho* Hall – NALIVKIN, pp. 8, 9, 71, pl. XI, figs 18a–d.  
 non 1939 – *Camarotoechia sappho* Hall – WILLARD, pp. 251, 289, 297, pl. 19, figs 22–24.  
 non 1939 – *Camarotoechia sappho* ? (Hall) – WILLARD, p. 290.  
 non 1942 – *Camarotoechia sappho* (Hall) – STUMM, p. 554, table, p. 556, pl. 84, fig. 11.  
 non 1943 – *Camarotoechia* cf. *sappho* (Hall) – REED, pp. 135–136.  
 1943 – *Camarotoechia sappho* (Hall) – REED, pp. 135, 136.  
 1944 – *Camarotoechia sappho* (Hall) – SHIMER & SHROCK, pl. 118, figs 57–60 (non fig. 61) (= pl. 54, figs 38–40, 42 in HALL, 1867).  
 e.p. 1944 – *Camarotoechia sappho* (Hall) – SHIMER & SHROCK, p. 311.  
 non 1947 – *Camarotoechia sappho* Hall – NALIVKIN, p. 29.  
 non 1956 – *Camarotoechia sappho* Hall – SIMORIN, pp. 21, 25.  
 e.p. 1956 – *Camarotoechia sappho* (Hall) – WANG, p. 143.  
 1958 – *Camarotoechia sappho* (Hall) (*Rhynchonella*) – STUMM & WRIGHT, pp. 92, 106, 115.  
 non 1959 – *Camarotoechia sappho* – WELLS, p. 36.  
 non 1959 – *Camarotoechia sappho* Hall – KRESTOVNIKOV in NAGIBINA & KRESTOVNIKOV, p. 8.  
 1959 – *Camarotoechia sappho* Hall – TORBAKOVA, p. 44.  
 non 1960 – *Camarotoechia sappho* – HOOVER, appendix, p. 138.  
 e.p. 1961 – *Camarotoechia sappho* (Hall) – FAGERSTROM, p. 27.  
 1963 – *Camarotoechia sappho* Hall – BUEHLER & TESMER, p. 46.  
 non 1963 – *Camarotoechia sappho* Hall – BUEHLER & TESMER, pp. 55, 58.  
 e.p. 1963 – *Camarotoechia sappho* Hall – NICOLAUS, p. 174.  
 non 1964 – *Camarotoechia sappho* (Hall) – BOBROV & MODZALEVSKAYA, p. 794.  
 non 1964 – *Camarotoechia sappho* (Hall) – TESMER, p. 45, table 1.  
 1964 – *Camarotoechia sappho* (Hall) – MANSPEIZER, pl. III, pl. IV.  
 e.p. 1964 – *Camarotoechia sappho* (Hall) – WANG et al., pp 357, 360.  
 non 1965 – *Camarotoechia sappho* Hall – MODZALEVSKAYA, pp. 8, 9.  
 1965 – *Cupularostrum sappho* (Hall) – GRIESEMER, p. 270.  
 1968 – *Camarotoechia sappho* Hall – RZHONSNITSKAYA, p. 260.  
 non 1969 – *Camarotoechia sappho* Hall – GUPTA, p. 9.  
 non 1972 – *Camarotoechia* sp., gr. *Sappho* Hall – ALAVI-NAINI, p. 63.  
 non 1973 – *Camarotoechia sappho* Hall – NALIVKIN et al., IX.  
 1977 – “*Camarotoechia*” *sappho* – GRASSO & WOLFF pl. 9, figs 7a,b.  
 e.p. 1981 – *Cupularostrum sappho* (HALL) – DUTRO, fig. 1, p. 69, p. 76, fig. 6, p. 77.  
 non 1985 – “*Camarotoechia*” *sappho* – MAMEDOV & RZHONSNITSKAYA, fig. 4, p. 143.  
 non 1986 – *Camarotoechia sappho* – MILLER, fig. 2, p. 61.  
 non 1993 – *Camarotoechia sappho* – BRETT, figs 12–17, p. 48 (= pl. 55, figs 47–52 in HALL, 1867).  
 1994 – *Cupularostrum sappho* – LINSLEY, p. 76.  
 non 1994 – *Cupularostrum sappho* – LINSLEY, p. 100.  
 non 1994 – *Cupularostrum sappho* – LINSLEY, pl. 82, fig. 29 (= pl. 54, fig. 33 in HALL, 1867).  
 1994 – *Cupularostrum sappho* – LINSLEY, pl. 82, figs 16–20, 27, 28, 30, 31 (= pl. 54, figs 34, 35, 37–43 in HALL, 1867).  
 non 1994 – *Cupularostrum sappho* – LINSLEY, pl. 82, figs 21–26 (= pl. 55, figs 47–52 in HALL, 1867).  
 non 1995 – «*Rhynchonella*» *sappho* – GARCÍA-ALCALDE, p. 25.  
 non 1995 – *Cupularostrum sappho* – BATT, fig. 7, p. 253.  
 1997 – *Cupularostrum sappho* – BRETT, p. 165.  
 non 1997 – *Camarotoechia sappho* – BRETT & VER STRAETEN (eds.), figs 12–17, p. 364 (= pl. 55, figs 47–52 in HALL, 1867).

#### Types, Locus typicus, Stratum typicum

The type series of *Rhynchonella sappho* described by HALL (1860, p. 87) and later figured with some slight additions [1867, pp. 340–341, pl. 54, figs 33–43 as *Rhynchonella (Stenocisma) sappho*] is made out of nine specimens. Eight of them are housed in the



Family Trigonirhynchiidae SCHMIDT, 1965		
SCHMIDT (1965, p. 2)	MCLAREN ( <i>in</i> SCHMIDT & MCLAREN, 1965, p. h559)	SAVAGE (1996, p. 252; 2002, p. 1052)
	small to medium-sized	
	uniplicate	
Rhynchonellacea von dreieckigem <i>bis</i> rundlichem Umriss		Rhynchotrematoidea with subtriangular outline
Naht kräftig gezackt	commisure serrate	anterior commissure <i>often</i> dentate, <i>sometimes</i> spinose
Sinus und Sattel <i>in der Regel</i> vorhanden	sulcus anf fold <i>commonly</i> developed	sulcus and fold variable
Rippen - kräftig - <i>meist einfach</i> - kantig <i>oder</i> gerundet-kantig - im Schnabel beginnend	costae - strong - simple - angular <i>or</i> subangular - extending from beak - <i>rarely</i> bifurcate	costae <i>often</i> extending from beaks, <i>but umbones commonly smooth</i>
Stirn und Schnabel <i>können</i> abgestutzt sein		
		delthyrium open <i>or</i> with disjunct-conjunct deltidial plates
In der Stielklappe <i>meist</i> gut ausgebildete Zahnstützen	dental plates present	dental plates present
In der Armklappe: - Septum - gut entwickeltes Septalium, das bei einigen Gattungen offen, bei anderen ± bedeckt ist.	median septum well-formed septalium that <i>may be open or wholly or partly covered by plate uniting outer hinge plates</i>	medium septum present septalium with or without cover plate
Schlossvorsatz fehlt		cardinal process absent

Fig. 1 — Original diagnosis of the family Trigonirhynchiidae SCHMIDT, 1965 parallel with those of the treatises (1965, 1996 + 2002). *Italics* indicate alternative characters.

American Museum of Natural History, New York [AMNH34623–34629 (formerly 4382), and 5214]; five specimens are figured (pl. 54, fig. 34–43 *in* HALL, 1867): four (figs 34–42) come from “limestone of the Marcellus shale” (HALL, 1860, p. 87) or from “calcareous layers in the Marcellus shale” (HALL, 1867, p. 341) “near Le Roy, Genesee County, New York” (HALL, 1860, p. 87; 1867, p. 341), the fifth one comes probably from near Le Roy, but WHITFIELD & HOVEY (1900, pp. 210–211) claim that it comes from Madison County. The ninth specimen (pl. 54, fig. 33) from the “Upper Helderberg limestones”, “assigned with some doubt” to the species, could not be located neither in the AMNH nor in any major American museum or scientific institution visited by the author. Six specimens (AMNH34630–34635) referring to a variety from Ohio, are not part of the type series, because they were first mentioned by HALL in 1867 (pl. 55, figs 47–52); two of them have been photographed (Pl.1, Fig. 34).

Lectotype, AMNH34626 (formerly 4382). Pl.1, Figs 1–5 = pl. 54, figs 38–42 *in* HALL, 1867. Limestone in the Marcellus shale, near Le Roy, Genesee County, New York. This specimen has been designated as the lectotype, because it was considered as a “characteristic specimen of the species” by HALL (1867) or a “typical example” by HALL & CLARKE (1893); figures of the lectotype have been reproduced many times in the literature (see Synonymy). Paralectotype A = pl. 54, fig. 33 *in* HALL, 1867. Upper Helderberg limestones”, “referred with some doubt to the species”. Paralectotypes B, AMNH34623 (formerly 4382), Pl.1, Figs 26–30 = pl. 54, figs 34, 35 *in* HALL, 1867, C, AMNH34624 (formerly 4382), Pl.1, Figs 21–25 = pl. 54, fig. 36 *in* HALL, 1867, D, AMNH34625 (formerly 4382), Pl.1, Figs 16–20 = pl. 54, fig. 37 *in* HALL, 1867, F, AMNH34627, G, AMNH34628, Pl.1, Figs 31–33, H, AMNH34629, Pl.1, Figs 6–10. Same locality and rock unit as the lectotype.



Paralectotype E, AMNH5214, Pl.1, Figs 11–15 = pl. 54, fig. 43 in HALL, 1867. Madison County, New York, according to WHITFIELD & HOVEY (1899, pp. 210–211).

Topotypes A–C, IRScNBa12215–12217. Batavia, Genesee County, western New York. Stafford Limestone. Collector: P. SARTENAER, 1953.

#### Other material

Outside the type series, with the exception of Paralectotype A that anyhow does not belong to the species, eight specimens collected by the author in 1953 from the Stafford limestone, Batavia, Genesee County, New York, have been studied. Many specimens housed in various museums and scientific institutions have also been examined.

#### Description

##### Remark

The external characters of this “beautiful and symmetrical” species, as HALL (1860, p. 87; 1867, p. 341) qualified it, were adequately described by HALL (1860, p. 87), but it is only some years later that HALL (1867, pp. 340–341, pl. 54, figs 33–43) figured the species and completed its original description. Subsequent descriptions by various authors deal only partly or not at all with the species as redefined in the present paper.

This refers only to specific characters in need of further elaboration.

Width of sulcus at front between 57 and 79 per cent (mostly between 61 and 74 per cent) of shell width. Top of tongue located lower than the highest part of the shell, i.e. between 16 and 25 per cent of shell thickness. Length of ventral interarea mostly between 24 and 35 per cent of shell width.

Measurements of the lectotype, seven paralectotypes, and two topotypes are given on Table 1; the lectotype and six paralectotypes have been photographed. Thickness of dorsal valve between 69 and 79 per cent (mostly 72 to 77 per cent) of shell thickness. Maximum thickness of dorsal valve, and thus of shell, between 12 and 43 per cent (mostly 24 to 30 per cent) of shell length anterior to the ventral beak. Maximum shell width between 45 and 65 per cent of shell length anterior to the ventral beak. Angle of the cardinal commissure between 115° and 128° (mostly between 120° and 128°).

The general costal formula in median, parietal, and lateral categories derived from at least 75 per cent of the specimens is:  $\frac{5 \text{ to } 6}{4 \text{ to } 5}$ ; 0 to  $\frac{1-0}{1-0}$ , and  $\frac{1-1}{1-1}$ ,  $\frac{6 \text{ to } 8}{7 \text{ to } 9}$ ,

ratios of costae of the lectotype, six paralectotypes, and two topotypes are given in the explanation of Plate 1. Parietal costae present in about half the specimens. Width of median costae at front varies between 2 and 2.5 mm.

Transverse serial sections of one specimen (topotype C IRScNBa12217) are shown on Text-figure 2; they are the first sections ever made from a specimen of the species.

#### Discussion of synonymy

Although rather complete, the synonymy list is by no means exhaustive. “e.p.” in front of some references means that they include also, but not only, the species. The negative part of the synonymy is explained in the section Stratigraphical Range and Geographical Distribution.

#### Comparisons

*Rhynchonella orbicularis* HALL, 1860 from the “Chemung Group” of SW New York, and also present in NW Pennsylvania and NE Ohio, is one of the species assigned to the genus *Cupularostrum* by GRIESEMER (1965, p. 270) (see above). DUTRO (1981, fig. 7, p. 79) mentions also *C. ? orbicularis*, although COOPER & DUTRO (1982, p. 71) prefer to put the species in the genus *Ripidiorhynchus* SARTENAER, 1966, an assignment already rejected by SARTENAER (1985, p. 323; 2001, p. 208). *Rhynchonella orbicularis*, which is “diagnostic of the Conewango Group” according to CHADWICK (1935, p. 328), is only mentioned here because CASTER (1930, pp. 152, 157, 158, 159 as *Camarotoechia sappho*) places the species described by HALL (1867) as *Rhynchonella (Stenocisma) orbicularis* and by HALL & CLARKE (1893) as *Camarotoechia orbicularis* into the synonymy of *Sapphicorhynchus sappho*. Although its appropriate generic assignment is not yet solved, *Rhynchonella orbicularis*, which is abundantly figured in the literature, differs from *Sapphicorhynchus sappho*, even in CASTER’s conception, by so many characters that there is no need to be more explicit.

Another species discussed in a PhD by HOLLAND (1959), and of which the generic status is also outstanding, is *Rhynchonella Allegania* WILLIAMS, 1887 known from SW New York and NW Pennsylvania as “highly characteristic” of Oswayo shales by BUTTS (1903, p. 99) and GOLDRING (1931, p. 428), “characteristic” of Oswayo beds by GOLDRING (1931, p. 426), a “Oswayo index fossil” by CASTER (1934, p. 95) and TESMER (1975, pp. 28, 66, 74, 79) or a “diagnostic species” of the Conewango






Fig. 2 — *Sapphicorhynchus sappho* (HALL, 1860). Topotype C, IRScNBa12217. Transverse serial sections, figures are in mm from ventral umbo. Measurements: length = 17.1 mm; width = 20.7 mm; thickness = 13.2 mm. Costal formula:  $\frac{6}{5}$ , 0;  $\frac{7}{8}$ .

Group by CHADWICK (1935, p. 329). The species also lent its name to a zone (RICKARD, 1975, pl. 3), an Assemblage Zone (DUTRO, 1981, fig. 2, p. 70, p. 81) or a fauna (CARTER & KAMMER, 1990, p. 94). Generally attributed to *Camarotoechia*, the species has been questionably assigned to *Sinotectirostrum* SARTENAER, 1970 by DUTRO (1981, fig. 2, p. 70, p. 81), and SARTENAER (1970, p. 13) indicated that the possibility of its inclusion in the genus *Centrorhynchus* SARTENAER, 1970 should be considered. In any case,

*Rhynchonella Allegania* is so strikingly different from *Sapphicorhynchus sappho* that any comparison is futile. The species is only alluded to here because CASTER (1930, p. 155) wrote that it "is in general appearance very similar to *Camarotoechia sappho*". This statement does not correspond to the reality, while the following observation by WILLIAMS (1887, p. 88) is acceptable: "it [*Rhynchonella Allegania*] approaches more nearly *R. orbicularis*".



in mm	Paralectotype E A5214	Lectotype 34626	Paralectotype H 34629	Paralectotype F 34627	Paralectotype D 34625	Paralectotype C 34624	Paralectotype B 34623	Paralectotype G 34628	Topotype A	Topotype B
l	21.3	20.2	18.4	16.3	16.1	15.7	14.4	14.1	16.9	15.6
w	27.4	27.6	25.6	18.5	22.6	19.8	17.1	18.4	19.5	19.4
lvv unrolled	34.5	32.5	130.5	23.9	26	25.2	20.7	22.2	28.5	26
t	21.3	17.8	17	12.6	15.1	15.4	10.8	13.1	13.8	15.5
tvv	4.8	5.5	3.8	3.5	3.2	3.4	2.9	3.1	2.7	3.8
tdv	16.5	12.3	13.2	9.1	11.9	12	7.9	10	11.1	11.7
l/w	0.78	0.73	0.72	0.88	0.71	0.79	0.84	0.77	0.87	0.80
t/w	0.78	0.64	0.66	0.68	0.67	0.75	0.63	0.71	0.71	0.80
t/l	1	0.88	0.92	0.77	0.94	0.95	0.75	0.93	0.82	0.99
angle of the cardinal commissure	128°	125°	126°	122°	124°	?	120°	120°	111°	115°

Table 1 — Measurements of the lectotype, seven paralectotypes, and two topotypes; the lectotype and paralectotypes B–E, G, H have been photographed. Abbreviations: l = length; w = width; t = thickness; vv = ventral valve; dv = dorsal valve.

*Stratigraphical range and geographical distribution*  
*Sapphicorhynchus sappho* comes only from the Stafford Limestone of western New York, and its time equivalent, the Mottville Sandstone and Limestone of central New York, both being the lowermost members of the Lower Givetian Skaneateles Formation. This approximately corresponds to DUTRO's (1981, fig. 1, p. 69) *Cupularostrum sappho* Occurrence Zone. All other references [Remark: It has been mentioned above (see Type species) that forms from the Centerfield Limestone, Ledyard Shale, Wanakah Shale, and Otisco Shale could belong to the genus, if not to the species] are indicative of other taxa and stratigraphical levels from the type area and outside New York State. The following concise review (New York references are in ascending stratigraphical order), which includes assignments with a query and varieties, runs parallel to the negative synonymy list.

*Western and Central New York*

*Upper Helderberg limestones*: HALL (1867 "referred with doubt to the species"), and by various authors referring to HALL, e.g. LESLEY (1898), KNOD (1908), FERUGLIO (1930), and LINSLEY (1994).

*Centerfield Limestone Member* (= lowermost member of the Middle Givetian Ludlowville Formation): SMITH (1935, p. 104).

*Otisco Shale Member* (= second from the base of the four members of the Ludlowville Formation in central

New York): SMITH (1935, p. 106).

*Wanakah Shale Member* (= third from the base of the four members of the Ludlowville Formation in western New York): GRABAU (1899, *Pleurodictyum* and *Trilobite* beds; 1899a,b, *Pleurodictyum* bed), BUEHLER & TESMER (1963), MILLER (1986, *Pleurodictyum* bed), and BATT (1995, *Trilobite* beds).  
*Tichenor Limestone Member* (= lower member of the Upper Givetian Moscow Formation in western New York): BUEHLER & TESMER (1963).

*Encrinal Limestone* (= base of the Moscow Formation): GRABAU (1899), SHIMER & GRABAU (1902), CLELAND (1903), RAYMOND (1904, *Encrinal limestone* and above), and STAUFFER (1915, p. 171).

*Portland Point Limestone* (in lower part of the Moscow Formation in central New York): SMITH (1935, p. 109).

*Windom Shale Member* (= upper member of the Moscow Formation): CLELAND (1903, *Cystodictya* Zone), RAYMOND (1904, p. 162, *Cystodictya* beds), and WELLS (1959).

*Tully Limestone*: WILLIAMS (1887), SMITH (1935, p. 110), COOPER & WILLIAMS (1935), and LINSLEY (1994, p. 100).

*Chemung Beds*: WILLIAMS (1884; 1887; in KINDLE, 1896), LESLEY (1890, p. 1060), and VON ENGELN (1932).

*Sherburne Member* (in middle part of the Genesee Formation): SMITH (1935, p. 111.)



*Ithaca Member* (in upper part of the Genesee Formation): CLARKE (1905).

*Canadaway Group*: CHADWICK (1935), and MANSPEIZER (1964).

*Conneaut Group*: MANSPEIZER (1964).

*Chadakoin Formation*: CASTER (1930).

*Ellicott Shale*: TESMER (1964).

*Conewango Group*: CASTER (1930), and CHADWICK (1935).

*Cattaraugus Formation*: TESMER (1964).

*Bradfordian, Owayo Formation, Waverly Group, Olean Conglomerate, Knapp Conglomerate, Salamanca Conglomerate*: LESLEY (1890, above and below the Salamanca conglomerate, "Chemung" beds), BUTTS (1903, Owayo shales, Knapp beds, Salamanca conglomerate, interval between the Wolf creek conglomerate and the top of the chocolate shales), and CHADWICK (1935, Owayo-Knapp of BUTTS).

*Carboniferous*: GIRTY (1899, p. 492: "As is well known, *Camarotoechia sappho* and *Spirifer subattenuatus* occur in both Devonian and Carboniferous deposits").

#### *North America outside of New York*

*Ontario*: In the upper Givetian Hungry Hollow and Widder Formations from the Thedford-Arkona region, SW Ontario, by WHITEAVES (1898), SHIMER & GRABAU (1902), RAYMOND (1904), STAUFFER (1915), and STUMM & WRIGHT (1958).

*Pennsylvania*: LESLEY [1889–1890, from the Upper Devonian of north-central Pennsylvania and from the Givetian Hamilton Group of central and north-central Pennsylvania (eight specimens of the collection identified by HALL)]; KINDLE in WILLIAMS & KINDLE (1905, from the Upper Devonian Chemung Group of north-central Pennsylvania); CASTER (1930, from the Cassadaga (Chadakoin) and Bradford (Conewango, Venango, Owayo) Stages; CASTER (1934, from the Strunian Bedford fauna of northwestern Pennsylvania); BUTTS in CHADWICK (1935, from the Conewango and Owayo of northwestern Pennsylvania; and WILLARD (1939, from the Catskill Group of northeastern Pennsylvania).

*Ohio*: HALL [1867; WHITFIELD & HOVEY (1899, pp. 210–211 did not mention HALL's variety in their table)], WILLIAMS (1887), HERRICK (1888), HERRICK in HALL & CLARKE (1893), GRABAU & SHIMER (1909), BRETT (1993), LINSLEY (1994), BRETT & VER STRAETEN (1997) from the Waverly Group of east-central Ohio; STAUFFER (1909 from the upper Eifelian Delaware Limestone of north-central Ohio);

PROSSER [1912, from the Famennian Erie (Chagrin) Shale of northeastern Ohio]; STUMM (1942, from the upper Givetian Prout Limestone of northeastern Ohio); HOOVER (1960).

*Indiana*: KINDLE (1899, pp. 13, 15, 16, 19) from the Lower Mississippian Riverside Sandstone of southern Indiana; 1901, from the uppermost Emsian-lower Eifelian Jeffersonville Limestone and the lower Givetian Sellersburg Beds of southern Indiana.

*Iowa*: LAUDON (1931) from the lower Famennian Sheffield Formation of north-central Iowa.

*Kentucky*: KINDLE (1899, from the Lower Mississippian Riverside Sandstone of central Kentucky; in WILLIAMS & KINDLE, 1905, from the Riverside Limestone and the Osagean Knobstone Formation of west-central Kentucky).

*Maryland*: PROSSER in PROSSER & KINDLE (1913) and in PROSSER *et al.* (1913) from the Givetian Hamilton Member of the Romney Formation of western Maryland.

*Missouri*: BRANSON & WILLIAMS (1922) from (probably) the Middle Devonian Grand Tower Limestone of southeastern Missouri.

*Montana*: GIRTY in KINDLE (1908) from the upper Carboniferous of southwestern Montana.

*Virginia*: KINDLE in WILLIAMS & KINDLE (1905) and WILLIAMS in WILLIAMS & KINDLE (1905) from the Kimberling Shale (Chemung fauna).

*West Virginia*: KINDLE in WILLIAMS & KINDLE (1905) and WILLIAMS in WILLIAMS & KINDLE (1905) from the Jennings blackish sandy shale (Chemung fauna) of southern West Virginia.

*Wyoming*: GIRTY (1899; in KINDLE, 1908) from the Mississippian Madison Limestone of the Yellowstone National Park.

#### *Outside of North America*

*Brazil*: THOMAS (1905) from the Devonian Maecurú Group of the States of Amazonas and Pará; KNOD (1908, referring to CLARKE, 1900); FERUGLIO (1930).

*England (Devon)*: REED (1943) from the Pilton Beds (Devonian/Carboniferous boundary beds) of North Devon.

*Iran*: STEPANOV in ALAVI-NAÏNI (1972) from the early Givetian part of the Bahram Formation.

*Kashmir*: GUPTA (1969) from the Couvinian.

*Kazakhstan*: NALIVKIN (1937, 1947) and SIMORIN (1956) from the Frasnian and Givetian.

*Mongolia*: BOBROV & MODZALEVSKAYA (1964) from the Middle Devonian.

*Russia: Arctic region*: NALIVKIN (1933) from the



Upper Devonian at the mouth of the Lena River; *Far East*: KRESTOVNIKOV in NAGIBINA & KRESTOVNIKOV (1959) and NALIVKIN *et al.* (1973) from the Givetian of the Amur River; *Transcaucasia*: MAMEDOV & RZHONSNITSKAYA (1985) from the late Givetian of Nakhichevan.

*Spain*: GARCÍA-ALCALDE (1995) from the Middle Devonian of the Cantabrian Zone.

## Acknowledgments

The author is grateful to the late Dr. N.D. Newell and to Dr. N. Eldredge, American Museum of Natural History, New York, for access to collections under their responsibility, long period loan, and permission to photograph types. In course of time the author had various fruitful conversations with the late Dr. G.A. Cooper, who allowed him to go through the collections of the National Museum of Natural History, Washington. Dr. G.J. Kloc, University of Rochester, kindly made some specimens of *Sapphicorhynchus sappho* available to the author. The author also wishes to express his appreciation to Prof. Dr. Michael R.W. Amler, Ludwig Maximilians-Universität München for critically reading the manuscript, and for helpful comments.

## References

- ALAVI-NAÏNI, M., 1972. Étude géologique de la région de Djam. *Report of the Geological Survey of Iran*, **23**: 288 pp.
- AMSDEN, T.W., 1974. Late Ordovician and early Silurian articulate brachiopods from Oklahoma, southwestern Illinois, and eastern Missouri. *Bulletin of the Oklahoma Geological Survey*, **119**: 154 pp.
- BARANOV, V.V., 1989. Novye devonskie rinkhonellidy Severo-Vostoka SSSR. *Paleontologicheskii Zhurnal*, **1**: 42–50.
- BATT, R.J., 1995. A test of a new technique illustrating faunal dominance trends: application to the “Trilobite Beds” interval of the Middle Devonian Wanakah Shale in western New York. *Lethaia*, **26** (3): 245–258.
- BAYLE, E., 1878. Explication de la carte géologique de la France, **4** (1), Fossiles principaux des terrains de la France, Atlas: 79 pls.
- BERDAN, J.M., 1964. The Helderberg Group and the position of the Silurian–Devonian boundary in North America. *Bulletin of the United States Geological Survey*, **1180-B**, 19 pp.
- BILLINGS, E., 1860–1861. On the Devonian fossils of Canada West (extracted from the Report of the Geological Survey of Canada for 1860, in preparation). *The Canadian Journal of Industry, Science, and Art*, **5** (27): 249–282, **6** (32, 33, 34): 138–148, 253–274, 329–363.
- BOBROV, V.A. & MODZALEVSKAYA, E.A., 1964. Novye dannye o srednem devone vostochnoy Mongolii. *Doklady Akademii Nauk SSSR*, **159** (4): 793–795.
- BRANSON, E.B., 1923. The Devonian of Missouri. Missouri Bureau of Geology and Mines, Second Series, **17** (1922): 1–129, 276–279.
- BRANSON, E.B. & WILLIAMS, J.S., 1922. Fauna of the Middle Devonian of southeastern Missouri. In: BRANSON, E.B., The Devonian of Missouri: 130–165. *Missouri Bureau of Geology and Mines*, **XVII**. Rolla, Missouri.
- BREIVEL, I.A. & BREIVEL, M.G., 1976. Genus *Losvia*. In: TYAZHEVA, A.P., ZHAVORONKOVA, R.A. & GARIFULLINA, A.A. (red.), Korally i brachiopody nizhnego devona Yuzhnogo Urala: 116–119. Nauka, Moskva.
- BRETT, C.E., 1993 (ed.). Field Trip Guidebook Central New York State. Sixty-first anniversary reunion of the Paleontological Research Institution, Ithaca, New York, August 13–15.
- BRETT, C.E., 1997. Day 1. Stop A–2 Oatka Creek, Leroy. In: BRETT, C.E. & VER STRAETEN, C.A. (eds.), Devonian cyclicity and sequence stratigraphy in New York State, Field Trip Guidebook, July 22–27, 1997: 165–166. University of Rochester.
- BRETT, C.E. & BAIRD, G.C., 1993. Middle Devonian stratigraphy, facies, and depositional environments of western New York State. In: BRETT, C.E. (ed.), Field Trip Guidebook Central New York State. Sixty-first anniversary reunion of the Paleontological Research Institution, Ithaca, New York, August 13–15: 5–11.
- BRETT, C.E. & VER STRAETEN, C.A. (eds.), 1997. Devonian cyclicity and sequence stratigraphy in New York State. Field Trip Guidebook, July 22–27, 1997: 369 pp. University of Rochester.
- BRICE, D., CARLS, P., COCKS, L.H.M., COPPER, P., GARCÍA-ALCALDE, J.L., GODEFROID, J. & RACHEBOEUF, P.R. (2000): Brachiopoda. In: BULTYNCK, P. (ed.), Subcommission on Devonian Stratigraphy, Fossil groups important for boundary definition. *Courier Forschungsinstitut Senckenberg*, **220**: 65–86.
- BUBLICHENKO, N.L., 1956. Nekotorye novye predstaviteli brachiopod devona i karbona Rudnogo Altaya i Sary-Arka. *Izvestiya Akademii Nauk Kazakhskoy SSR, seriya geologicheskaya*, **23**: 93–104.
- BUEHLER, E.J. & TESMER, I.H., 1963. Geology of Erie County, New York. *Bulletin of the Buffalo Society of Natural Sciences*, **21** (3).
- BUTTS, C., 1903. Fossil faunas of the Olean quadrangle. Appendix 3, Report of the State Paleontologist 1902: 990–995. *Bulletin of the New York State Museum*, **69**, Paleontology 9.



- CARTER, J.L. & KAMMER, T.W., 1990. Late Devonian and early Carboniferous brachiopods (Brachiopoda, Articulata) from the Price Formation of West Virginia and adjacent areas of Pennsylvania and Maryland. *Annals of the Carnegie Museum*, **59** (2): 77–103.
- CASTER, K.E., 1930. Higher fossil faunas of the Upper Allegheny. *Bulletins of American Paleontology*, **15** (58): 113 p.
- CASTER, K.E., 1934. The stratigraphy and paleontology of northwestern Pennsylvania, Part I: Stratigraphy. *Bulletins of American Paleontology*, **21** (71): 185 p.
- CHADWICK, G.H., 1935. Faunal differentiation in the Upper Devonian. *Bulletin of the Geological Society of America*, **46** (2): 305–342.
- CLARKE, J.M., 1901. Limestones of central and western New York interbedded with bituminous shales of the Marcellus stage with notes on the nature and the origin of their faunas. *Bulletin of the New York State Museum*, **50**, *Paleontologic Papers*, **2**: 115–138.
- CLARKE, J.M., 1905. Ithaca fauna of central New York. *Bulletin of the New York State Museum*, **82**, *Paleontology*, **12**: 53–65, 67–70.
- CLARKE, J.M. & SWARTZ, C.K., 1913. Systematic paleontology of the Upper Devonian deposits of Maryland. Brachiopods. Middle and Upper Devonian volume: 546–605. Baltimore, Maryland.
- CLELAND, H.F., 1903. A study of the fauna of the Hamilton formation of the Cayuga Lake section in central New York. *Bulletin of the United States Geological Survey*, **206**: 112 pp.
- CLELAND, H.F., 1911. The fossils and stratigraphy of the Middle Devonian of Wisconsin. *Bulletin of the Wisconsin Geological and Natural History Survey*, **21**, Scientific series, 222 pp.
- CONRAD, T.A., 1841. Fifth Annual Report on the Palaeontology of the State of New-York. Annual Report of the New-York Geological Survey: **5**. State of New-York, n°150 in Assembly, February 17: 25–57.
- COOPER, G.A., 1929. Stratigraphy of the Hamilton Group of New York. PhD, Yale University.
- COOPER, G.A., 1942. New genera of North American brachiopods. *Journal of the Washington Academy of Sciences*, **32** (8): 228–235.
- COOPER, G.A., 1956. Chazy and related brachiopods. *Smithsonian Miscellaneous Collections*, **127** (1): 1–1024, (2): 1025–1245. Publication 4253.
- COOPER, G.A. & WILLIAMS, J.S., 1935. Tully formation of New York. *Bulletin of the Geological Society of America*, **46** (5): 781–868.
- COOPER, G.A. & DUTRO, J.T., Jr., 1982. Devonian brachiopods of New Mexico. *Bulletins of American Paleontology*, **82** & **83** (315), 215 pp.
- DROT, J. & WESTBROEK, P., 1966. *Iberirhynchia santaluciensis* nouveau *Rhynchonellacea* du Dévonien de Léon (Espagne). *Leidse Geologische Mededelingen*, **38**: 165–172.
- DUTRO, J.T., Jr., 1981. Devonian brachiopod biostratigraphy. In: OLIVER, W.A., Jr. & KLAPPER, G. (eds.), Devonian biostratigraphy of New York. Guidebook (Part 1 = text, Part 2 = stop descriptions). International Subcommittee on Devonian Stratigraphy, July 1981: 67–82. Washington.
- FAGERSTROM, J.A., 1961. The fauna of the Middle Devonian Formosa Reef Limestone of southwestern Ontario. *Journal of Paleontology*, **35** (1): 1–48.
- FERUGLIO, E., 1930. Fossili devonici del Quemado (San Pedro de Jujuy) nella regione subandina dell' Argentina settentrionale. *Giornale di Geologia*, serie 2a, **5**: 71–97.
- FISCHER DE WALDHEIM, G., 1809. Notice sur des fossiles du Gouvernement de Moscou, 1, Sur les coquilles fossiles dites Térébratules. – Invitation à la séance publique de la Société Impériale des Naturalistes de Moscou, le 26 octobre 1809: 35 p., 3 pls. Moscou.
- GARCÍA-ALCALDE, J. L., 1995. L'évolution paléogéographique pré-varisque de la Zone Cantabrique septentrionale (Espagne). *Revista Española de Paleontología*, **10** (1): 9–29.
- GARCÍA-ALCALDE, J.L., 1998. Braquiópodos rinconélidos del Lochkoviense (Devónico Inferior) de la Cordillera Cantábrica (Norte de España). *Geobios*, **31** (6): 767–789.
- GIRTY, G.H., 1899. Devonian and Carboniferous fossils. Geology of the Yellowstone National Park. Part II: Description, Geology, Petrography, and Paleontology. *Monographs of the United States Geological Survey*, **32** (II): 479–599.
- GOLDRING, W., 1931. Handbook of paleontology for beginners and amateurs, Part 2: The formations. *Handbooks*, **10**. New York State Museum.
- GRABAU, A.W., 1898–1899. Geology and palaeontology of 18 Mile Creek and the lake shore sections of Erie County, N.Y., Part 1: Geology, Part 2: Palaeontology. A hand book for the use of students and amateurs. *Buffalo Society of Natural Sciences*, **6** (1 = Pt. 1, 2–4 = Pt. 2).
- GRABAU, A.W., 1899a. The faunas of the Hamilton Group of Eighteen-mile Creek and vicinity, in western New York. 50th Annual Report of the Regents 1896, v. 2: Report of the State Geologists and Field Assistants: 227–339. New York State Museum, Albany.
- GRABAU, A.W., 1899b. The faunas of the Hamilton Group of Eighteen-mile Creek and vicinity, in western New York.



- Sixteenth Annual Report of the State Geologist for the year 1896: 227-440.
- GRABAU, A.W., 1931-1933. Devonian Brachiopoda of China, 1: Devonian Brachiopoda from Yunnan and other districts in South China. *Palaeontologica Sinica*, **3** (3), 545 pp.
- GRABAU, A.W. & SHIMER, H.W., 1909. North American Index Fossils, Invertebrates, **1**: 853 pp. A.G. Seiler & Company, New York.
- GRASSO, T.X. & WOLFF, M.F., 1977. Paleoenvironments of the Marcellus and Lower Skaneateles Formations of the Otsego County region (Middle Devonian). Guidebook to Field Excursions conducted at the 49th annual meeting of the New York State Geological Association, September 1977, Trip A-3: 1-39, 42-50.
- GRIESEMER, A.D., 1965. Middle Devonian brachiopods of southeastern Wisconsin. *The Ohio Journal of Science*, **65** (5): 241-295.
- GUPTA, V.J., 1969. Palaeozoic stratigraphy of the area southeast of Srinagar, Kashmir. *Research Bulletin of the Panjab University*, New series (Sciences), **20** (I): 1-14.
- HALL, J., 1843. Geology of New York. Part IV. Survey of the Fourth Geological District: 683 pp. Albany.
- HALL, J., 1860. Contributions to palaeontology 1858 and 1859. 13th Annual Report of the Regents of the University of the State of New York: 55-125.
- HALL, J., 1867. Natural History of New York. Part VI: Palaeontology. Volume IV, Part I containing descriptions and figures of the fossil brachiopods of the upper Helderberg, Hamilton, Portage and Chemung groups, 428 pp. Geological Survey of the State of New York, Albany.
- HALL, J. & CLARKE, J.M. (1893-1894): Natural History of New York, Part VI, Palaeontology, vol. VIII: An introduction to the study of the genera of Palaeozoic Brachiopoda, Part II: 394 pp., 84 pls. Geological Survey of the State of New York, Albany.
- HAVLÍČEK, V., 1959. Rhynchonellacea im böhmischen älteren Paläozoikum (Brachiopoda). *Věstník Ústředního Ústavu Geologického*, **34** (1): 78-82.
- HAVLÍČEK, V., 1982. New genera of rhynchonellid and camerellid brachiopods in the Silurian of Bohemia. *Věstník Ústředního Ústavu Geologického*, **57** (6): 365-372.
- HAVLÍČEK, V. & ŠTORCH, P., 1990. Silurian brachiopods and benthic communities in the Prague Basin (Czechoslovakia). *Rozpravy Ústředního Ústavu Geologického*, **48**: 275 p.
- HERRICK, C.L., 1887-1888. A sketch of the geological history of Licking County accompanying an illustrated catalogue of Carboniferous fossils from Flint Ridge, Ohio. *Bulletin of the Scientific Laboratories of Denison University*, **II** (1), 1887: 5-68; **II** (2), 1887: 144-148; **III** (1), 1888: 13-110.
- HOLLABD, F.D., Jr., 1959. The stratigraphy and paleontology of northwestern Pennsylvania, Part II: Paleontology, Section B: The Brachiopoda of the Oswayo and Knapp Formations. PhD, University of Cincinnati, Ohio.
- HOOVER, K.V., 1960. Devonian-Mississippian shale sequence in Ohio. *Information Circular of the Geological Survey of Ohio*, **27**: 154 p.
- HOTZ, E.-E., KRÖMMELBEIN, K., KRAÜSEL, W. & STRUVE, W., 1955. Zur Geologie der Eifelkalkmulden. *Beihefte zum Geologisches Jahrbuch*, **17**, 204 pp.
- JIN, J., CALDWELL, W.G.E. & NORFORD, B.S., 1993. Early Silurian brachiopods and biostratigraphy of the Hudson Bay Lowlands, Manitoba, Ontario, and Quebec. *Bulletin of the Geological Survey of Canada*: **457**, 221 pp.
- JOHNSON, J.G. & BOUCOT, A.J., 1973. Devonian brachiopods. In: HALLAM, A. (ed.), Atlas of Palaeobiogeography: 89-96. Elsevier Scientific Publishing Company. Amsterdam, London, New York.
- JOHNSON, J.G. & PERRY, D.G., 1976. Middle Devonian brachiopods from the Bird Fiord Formation of Bathurst Island, Arctic Canada. *Canadian Journal of Earth Sciences*, **13** (5): 615-635.
- KINDLE, E.M., 1896. The relation of the fauna of the Ithaca group to the faunas of the Portage and the Chemung. *Bulletins of American Paleontology*, **2** (6): 54 pp.
- KINDLE, E.M., 1899. The Devonian and Lower Carboniferous faunas of southern Indiana and central Kentucky. *Bulletins of American Paleontology*, **3** (12): 111 pp.
- KINDLE, E.M., 1901. The Devonian fossils and stratigraphy of Indiana. *Annual Report of the State Geologist*, 1900, **25**: 529-758.
- KINDLE, E.M., 1908. The fauna and stratigraphy of the Jefferson limestone in the Northern Rocky Mountain region. *Bulletin of American Paleontology*, **4** (20): 79-117, pls 14-17; Ithaca.
- KNOD, R., 1908. Devonische Faunen Boliviens. In: STEINMANN, G., Beiträge zur Geologie und Paläontologie von Südamerika. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie*, **25** (Beilage Band): 493-600.
- LA TOUCHE, T.H.D. (1913): Geology of the northern Shan States. *Memoirs of the Geological Survey of India*, **39** (2): 379 pp. + i-iv, i-xli. Calcutta.
- LAUDON, L.R., 1931. A *Spirifer disjunctus* fauna in Iowa. *Proceedings of the Iowa Academy of Sciences*, **37** (1930): 251-255.
- LAURIE, J.R., 1991. Articulate brachiopods from the Ordovician and lower Silurian of Tasmania. In: JELL, P.A. (ed.), Australian Ordovician Brachiopod Studies. *Memoirs of the Association of Australian Palaeontologists*, **11**: 1-106.



- LENZ, A.C. (1971): *Werneckeella*, a new Lower Devonian rhynchonellid brachiopod, Royal Creek, Yukon Territory. *Journal of Paleontology*, **45** (5): 844–848.
- LESLEY, J.P., 1889–1890. A dictionary of the fossils of Pennsylvania and neighboring states named in the reports and catalogues of the Survey. *Reports of the Geological Survey of Pennsylvania*, **P4**, 3 volumes: 1283 pp. Harrisburg.
- LINSLEY, D.M., 1994. Devonian Paleontology of New York. *Special Publications of the Paleontological Research Institution*, **21**: 472 pp. Ithaca.
- LYASHENKO, A.I., 1973. Brachiopody i stratigrafiya nizhnefranskikh otlozheniy yuzhnogo Timana i Volgo-Uralskoy neftegazonosnoy provintsii. *Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Nefryanogo Instituta* (VNIGNI), **134**, 279 pp.
- MAMEDOV, A.B. & RZHONSNITSKAYA, M.A., 1985. Devonian of the South Transcaucasus. Zonal subdivision, boundaries of series and stages, correlation. In: ZIEGLER, W. & WERNER, R. (eds.), *Devonian Series Boundaries – Results of world-wide Studies. Courier Forschungsinstitut Senckenberg*, **75**: 135–156.
- MANSPEIZER, W., 1964. A study of the stratigraphy, paleontology, petrology, and geological history of the Canadaway and Conneaut Groups in Allegany County, New York. PhD thesis: 354 p., Rutgers State University.
- MERGL M., MASSA, D. & PLAUCHUT, B., 2001. Devonian and Carboniferous brachiopods and bivalves of the Djado Sub-Basin (North Niger, SW Libya). In: FRÝDA, J., BLODGETT, R.B. & MERGL, M. (eds.): 169–188. *Journal of the Czech Geological Society*, **46** (3–4, HAVLÍČEK Volume).
- MILLER, K.B., 1986. Depositional environments and sequences, “*Pleurodictyum* Zone”, Ludlowville Formation of western New York. In: BRETT, C.E. (ed.), *Dynamic stratigraphy and depositional environments of the Hamilton Group (Middle Devonian) in New York State, Part I. Bulletin of the New York State Museum*, **457**: 57–77.
- MODZALEVSKAYA, E.A., 1965. Paleontologicheskoe obosnovanie stratigrafii paleozoya Dal'nego Vostoka i Zabaykalya. *Biostratigraficheskiy Sbornik*, **1**. *Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Geologicheskogo Instituta* (VSEGEI), **115**: 3–12.
- NAGIBINA, M.S. & KRESTOVNIKOV, V.N., 1959. Skhema stratigrafii paleozoyskikh otlozheniy okrestnostey goroda Zei. *Geologiya i Razvedka*, **11**: 3–19.
- NALIVKIN, D.V., 1933. Devon Kharaulakhskikh gor, ostrova Kotel'nogo i Indigirki. *Trudy Arkticheskogo Instituta*, **12**: 203–212.
- NALIVKIN, D.V., 1937. Brachiopoda of the Upper and Middle Devonian and Lower Carboniferous of Northeastern Kazakhstan. *Trudy Tsentral'nogo Nauchno-Issledovatel'skogo Geologo-Razvedochnogo Instituta* (TsNIGRI), **99**: 200 pp.
- NALIVKIN, D.V., 1947. Klass *Brachiopoda*. Brachiopody. In: NALIVKIN, D.V. (ed.), *Atlas rukovodyashchikh form iskopaemykh faun SSSR*, tom III, *Devonskaya Sistema*: 63–134. Vsesoyuzny Geologicheskii Institut, Ministerstvo Geologii SSSR. Gosgeolizdat, Moskva, Leningrad.
- NALIVKIN, D.V., RZHONSNITSKAYA, M.A. & MARKOVSKIY, B.P.†, 1973. *Devonskaya Sistema, Stratigrafiya SSSR*, **1**: 520 p., **2**: 376 p. Izdatel'stvo “Nedra”, Moskva.
- NICOLAUS, H.-J., 1963. Zur Stratigraphie und Fauna der *crenistria*-Zone im Kulm des Rheinischen Schiefergebirges. *Beihefte zum Geologischen Jahrbuch*, **53**: 246 p.
- NIKIFOROVA, O.I. & ANDREEVA, O.N., 1961. Biostratigrafiya paleozoya Sibirskoy Platformy. Vypusk 1: Stratigrafiya ordovika i silura Sibirskoy Platformy i ee paleontologicheskoe obosnovanie (brachiopody). *Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Geologicheskogo Instituta* (VSEGEI): **56**.
- NORRIS, A.W., 1979. Devonian in the Western Hemisphere. In: MOORE, R.C. (founder), *Treatise on Invertebrate Paleontology, Part A, Introduction, Fossilization (Taphonomy), Biogeography and Biostratigraphy*: A218–A253. Boulder, Colorado and Lawrence, Kansas.
- OEHLERT, D., 1884. Études sur quelques Brachiopodes dévoniens. *Bulletin de la Société géologique de France*, **12** (1883–1884): 411–441.
- PROSSER, C.S., 1912. The Devonian and Mississippian formations of northeastern Ohio. *Bulletin of the Geological Survey of Ohio*, **15**: 340 p.
- PROSSER, C.S. & KINDLE, E.M., 1913. Systematic paleontology of the Middle Devonian deposits of Maryland. In: *Middle and Upper Devonian volume, Maryland Geological Survey*: 124–213. Baltimore.
- PROSSER, C.S., KINDLE, E.M. & SWARTZ, C.K., 1913. The Middle Devonian deposits of Maryland. In: *Middle and Upper Devonian volume*: 23–114. Baltimore.
- RACHEBOEUF, P.R., GIRARD, C., LETHIERS, F., DERYCKE, C., HERRERA, Z.A. & TROMPETTE, R., 2001. Evidence for Givetian stage in the Mauritanian Adrar (West Africa): biostratigraphical data and palaeogeographic implications. *Newsletters on Stratigraphy*, **38** (2/3): 141–162.
- RAYMOND, P.E., 1904. The *Tropidoleptus* fauna at Canandaigua Lake, New York, with the ontogeny of twenty species. *Annals of the Carnegie Museum*, **3** (1): 79–177.
- REED, F.R.C., 1943. Notes on certain Upper Devonian brachiopods figured by Whidborne. *Geological Magazine*, **80** (2 = Part 1: 69–78; 3 = Part 2: 95–106; 4 = Part 3: 132–138).



- RICKARD, L.V., 1975. Correlation of the Silurian and Devonian rocks in New York State. *Map and Chart Series*, 24. New York State Museum and Science Service. Albany.
- ROEMER, C.F., 1844. Das Rheinische Uebergangsgebirge, Eine palaeontologisch-geognostische Darstellung. Verlag der Hahn'schen Hofbuchhandlung, 96 pp.
- ROZMAN, Kh.S., 1969. Pozdneordoviskie brakhiopody Sibirskoy Platformy. *Paleontologicheskij Zhurnal*, 3: 86–108.
- RZHONSNITSKAYA, M.A., 1956. Tip Brachiopoda. In: KIPARISOVA, L.D., MARKOVSKIY, B.P. & RADCHENKO, G.P. (red.), Materialy po Paleontologii. Novye semeistva i rody. *Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Geologicheskogo Instituta (VSEGEI)*, novaya seriya, Paleontologiya, 12.
- RZHONSNITSKAYA, M.A., 1968. Biostratigrafiya devona okrain Kuznetskogo basseyna, tom I, Stratigrafiya. *Trudy Vsesoyuznogo Nauchno – Issledovatel'skogo Geologicheskogo Instituta (VSEGEI)*, 288 pp. "Nedra", Leningrad.
- SARTENAER, P., 1961. Late Upper Devonian (Famennian) rhynchonelloid brachiopods. *Bulletin de l'Institut royal des Sciences naturelles de Belgique*, 37 (24): 10 pp.
- SARTENAER, P., 1961. Etude nouvelle, en deux parties, du genre *Camarotoechia* HALL ET CLARKE, 1893. Deuxième partie: *Cupularostrum recticostatum* n. gen., n. sp. *Bulletin de l'Institut royal des Sciences naturelles de Belgique*, 37 (25): 15 pp.
- SARTENAER, P., 1966. *Ripidiorhynchus*, nouveau genre de Brachiopode Rhynchonellide du Frasnien. *Bulletin de l'Institut royal des Sciences naturelles de Belgique*, 42 (30): 1–15.
- SARTENAER, P., 1970. Nouveaux genres Rhynchonellides (Brachiopodes) du Paléozoïque. *Bulletin de l'Institut royal des Sciences naturelles de Belgique*, 46 (32): 32 pp.
- SARTENAER, P., 1985. In: ZIEGLER, W. & WERNER, R. (eds), Devonian Series Boundaries – Results of world-wide Studies. The biostratigraphical significance of rhynchonellid genera at the Givetian–Frasnian Boundary. *Courier Forschungsinstitut Senckenberg*, 75: 311–317. Frankfurt am Main.
- SARTENAER, P., 2001. Revision of the rhynchonellid brachiopod genus *Ripidiorhynchus* SARTENAER. *Geologica Belgica*, 3 (2000) (3–4): 191–213.
- SARTENAER, P., 2003. *Tullypothyridina*, new late Givetian rhynchonellid (brachiopod) genus. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, 73: 29–51.
- SARTENAER, P., 2004. *Cherryvalleyrostrum*, a new late Eifelian rhynchonellid (brachiopod) genus from North America. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, 74: 5–18.
- SAVAGE, N.M., 1996. Classification of Paleozoic rhynchonellid brachiopods. In: COPPER, P. & JIN, J. (eds), Brachiopods, Proceedings of the Third International Brachiopod Congress, Sudbury, Ontario, Canada, 2–5 September 1995, 249–260, 11 tables. A.A. Balkema, Rotterdam, Brookfield.
- SAVAGE, N.M., 2002. Superfamily Rhynchotrematoidea. In: MOORE, R.C. (founder), KAESLER, R.L. (ed.), Brachiopoda, revised. Part H of Treatise on Invertebrate Paleontology, 4, Rhynchonelliformea (part): 1047–1091. The Geological Society of America, Boulder, and The University of Kansas Press, Lawrence.
- SAVAGE, N.M., PERRY, D.G. & BOUCOT, A.J., 1979. A quantitative analysis of Lower Devonian brachiopod distribution. In: GRAY, J. & BOUCOT, A.J. (eds.), Historical, Biogeography, Plate Tectonics and the Changing Environment: 169–200. Oregon State University.
- SAVAGE, T.E., 1929. Tully fauna at the base of the black shale in east-central Kentucky. Proceedings of the forty-first annual meeting of the Geological Society of America and of the twentieth annual meeting of the Paleontological Society, held at New York City, December 1928. *Bulletin of the Geological Society of America*, 40 (1): 112, 249.
- SCHMIDT, H., 1941. Die mitteldevonischen Rhynchonelliden der Eifel. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft*, 459: 79 p.
- SCHMIDT, H., 1965. Neue Befunde an paläozoischen Rhynchonellacea (Brachiopoda). *Senckenbergiana lethaea*, 46 (1): 1–25.
- SCHMIDT, H. & McLAREN, D.J., 1965. Paleozoic Rhynchonellacea. In: MOORE, R.C. (ed.), Brachiopoda, Part H of Treatise on Invertebrate Paleontology: H552–597. – The Geological Society of America, Boulder, and The University of Kansas Press, Lawrence.
- SCHNUR, J., 1851. Die Brachiopoden aus dem Uebergangsgebirge der Eifel. Programm der vereinigten höhern Bürger- u. Provinzial-Gewerbeschule zu Trier, Schuljahr 1850/51: 2–16. Trier.
- SCHNUR, J. 1853. Zusammenstellung und Beschreibung sämtlicher im Übergangsgebirge der Eifel vorkommenden Brachiopoden nebst Abbildungen derselben. *Palaeontographica*, 3: 169–247.
- SCHUCHERT, C., 1903. On the faunal provinces of the Middle Devonian of America and the Devonian coral sub-provinces of Russia, with two paleogeographic maps. *American Geologist*, 32: 137–143, 149–150.
- SCHUCHERT, C., 1929. Classification of brachiopod genera, fossil and recent. In: SCHUCHERT, C. & LE VENE, C.M., Brachiopoda (Genorum and Genotyporum Index



- et Bibliographia). In: POMPECKI, J.F. (ed.), *Fossilium Catalogus I: Animalia*, **42**: 10–25. W. Junk, Berlin.
- SHIMER, H.W. & GRABAU, A.W., 1902. Hamilton group of Thedford, Ontario. *Bulletin of the Geological Society of America*, **13**: 149–186.
- SHIMER, H.W. & SHROCK, R.R., 1944. Index fossils of North America: 837 pp. + i–ix. John Wiley & Sons, New York. Chapman & Hall, London.
- SIMORIN, A.M., 1956. Stratigrafiya i brakhiopody Karagandinskogo basseyna. Institut Geologicheskikh Nauk. Akademiya Nauk Kazakhskoy SSR: 300 pp. Alma-Ata.
- SMITH, B., 1935. Geology and mineral resources of the Skaneateles quadrangle. *Bulletin of the New York State Museum*, **300**: 120 pp.
- SPIESTERSBACH, J. & FUCHS, A., 1909. Die Fauna der Remscheider Schichten. *Abhandlungen der Königlich Preussischen Geologischen Landesanstalt*, N.F., **58**: 1–81.
- STAINBROOK, M.A., 1942. Brachiopoda of the Cedar Valley beds of Iowa – Inarticulata, Rhynchonellacea and Rostrospiracea. *Journal of Paleontology*, **16** (5): 604–619.
- STAINBROOK, M.A., 1945. Brachiopoda of the Independence Shale of Iowa. *Memoirs of the Geological Society of America*, **14**, 74 pp.
- STAUFFER, C.R., 1909. The Middle Devonian of Ohio. *Bulletin of the Geological Survey of Ohio*, **10**: 204 pp.
- STAUFFER, C.R., 1915. The Devonian of southwestern Ontario. *Memoirs of the Geological Survey of Canada*, **34** (Geological Series, 63): 341 pp.
- STUMM, E.C., 1942. Fauna and stratigraphic relations of the Prout limestone and Plum Brook shale of northern Ohio. *Journal of Paleontology*, **16** (5): 549–563.
- STUMM, E.C. & WRIGHT, J.D., 1958. Check list of fossil invertebrates described from the Middle Devonian rocks of the Thedford–Arkona region of southwestern Ontario. *Contributions of the Museum of Paleontology of the University of Michigan*, **14** (7): 81–132.
- TEMPLE, J.T., 1965. Upper Ordovician brachiopods from Poland and Britain. *Acta Palaeontologica Polonica*, **10** (3): 379–427.
- TESMER, I.H., 1964. Geology of Chautauqua County, New York. Part I: Stratigraphy and paleontology (Upper Devonian). *Bulletin of the State Museum*, **391** (1963).
- TESMER, I.H., 1975. Geology of Cattaraugus County, New York. *Bulletin of the Buffalo Society of Natural Sciences*, **27**.
- THOMAS, I., 1905. Neue Beiträge zur Kenntnis der devonischen Fauna Argentinien. *Zeitschrift der Deutschen geologischen Gesellschaft*, **57** (2): 233–290.
- TORBAKOVA, A.F., 1959. Opisanie fauny. Tip Brachiopoda. Nadsemeystvo Rhynchonellacea. In: KHODALEVICH, A.N., BREIVEL, I.A., BREIVEL, M.G., VAGANOVA, T.I., TORBAKOVA, A.F. & YANET, F.E., Brakhiopody i korally iz eifelskikh boksitososnykh otlojeniy vostochnogo sklona Srednego i Severnogo Urala: 42–52. Ministerstvo Geologii i Okhrany Nedr SSSR, Ural'skoe Geologicheskoe Upravlenie.
- ULRICH, E.O. & COOPER, G.A., 1942. New genera of Ordovician brachiopods. *Journal of Paleontology*, **16** (5): 620–626.
- VANUXEM, L., 1842. Survey of the third geological district. In: Natural History of New York, Part IV: Geology, Pt. III: 306 pp. Albany.
- VON ENGELN, O.D., 1932. The Finger Lake region. In: NEWLAND, D.H. (dir.), The Paleozoic stratigraphy of New York, XVI International Geological Congress, United States, Guidebook 4, Excursion A–4: 39–69.
- WANG, Y., 1956. Some new brachiopods from the Yukiang Formation of southern Kwangsi Province. *Acta Palaeontologica Sinica*, **IV** (2): 137–162. (in Chinese with English Summary).
- WANG, Y. et al., 1964. Brachiopod fossils of China, 2 vol., 777 pp., 136 pl. Scientific Press, Beijing. (in Chinese).
- WEDEKIND, R., 1924. Das Mitteldevon der Eifel. I. Die Tetrakorallen des unteren Mitteldevon. *Schriften der Naturwissenschaftlichen Gesellschaft Marburg*, **14** (3): 1–93.
- WELLS, J.W., 1959. Geology of the Cayuga Lake Basin. A guide for the 31st annual field meeting of the New York State Geological Association prepared by staff and students of the Department of Geology, Cornell University. Second (revised) edition. Ithaca, New York.
- WESTBROEK, P., 1964. Systématique et importance stratigraphique des Rhynchonelles du Calcaire de Crémènes (Dévonien Supérieur, Province de León, Espagne). *Leidsche Geologische Mededelingen*, **30**: 243–252.
- WHITEAVES, J.F., 1898. On some additional or imperfectly understood fossils from the Hamilton formation of Ontario, with a revised list of the species therefrom. *Contributions to Canadian Palaeontology*, **1** (5): 361–363, 386–388, 392–394, 412, 415–416.
- WHITFIELD, R.P. & HOVEY, E.O., 1899–1900. Catalogue of the types and figured specimens in the palaeontological collection of the Geological Department, American Museum of Natural History. *Bulletin of the American Museum of Natural History*, **11** (2, 1899: 73–188; 3, 1900: 189–356).
- WILLARD, B., 1939. The Devonian of Pennsylvania.



Middle and Upper Devonian. *Bulletin of the Pennsylvania Geological Survey*, **G19**: 1–28, 131–381.

WILLIAMS, H.S., 1884. On the fossil faunas of the Upper Devonian along the meridian of 76°30' from Tompkins County, New York, to Bradford County, Pennsylvania. *Bulletin of the United States Geological Survey*, **1** (3): 1–36.

WILLIAMS, H.S., 1887. On the fossil faunas of the Upper Devonian. The Genesee section, New York. *Bulletin of the United States Geological Survey*, **41**: 31–103.

WILLIAMS, H.S. & KINDLE, E.M., 1905. Contributions to Devonian paleontology 1903, Part II, Fossil faunas of Devonian sections in central and northern Pennsylvania. *Bulletin of the United States Geological Survey*, **244**: 59–144.

WILLIAMS, S.G., 1887. The Tully limestone, its distribution and its known fossils [communicated to the Report of the State Geologist for 1886]. *Sixth Annual Report of the State Geologist for the year 1886*: 13–29.

WOOD, E., 1901. Marcellus (Stafford) limestones of Lancaster, Erie Co. N.Y. *Bulletin of the New York State Museum*, **49** (Paleontological Papers **2**): 139–181.

Paul SARTENAER  
Department of Palaeontology  
Royal Belgian Institute of Natural Sciences  
rue Vautier 29, B-1000 Brussels, Belgium

Typescript received: August 31, 2006

Revised typescript received: May 15, 2007



### Explanation of Plate 1

All figures are natural size

- Figs 1-5 — *Sapphicorhynchus sappho* (HALL, 1860). Lectotype, AMNH34626 = pl. 54, figs 38–42 in HALL, 1867. Dorsal, ventral, anterior, posterior, and lateral views. Costal formula:  $\frac{6}{5}$ ;  $\frac{1-0}{1-0}$ ;  $\frac{8}{8}$  and  $\frac{9}{9}$ .
- Figs 6-10 — *Sapphicorhynchus sappho* (HALL, 1860). Paralectotype H, AMNH34629. Dorsal, ventral, anterior, posterior, and lateral views. Costal formula:  $\frac{6}{5}$ ; 0;  $\frac{7}{8}$ .
- Figs 11-15 — *Sapphicorhynchus sappho* (HALL, 1860). Paralectotype E, AMNH5214 = pl. 54, fig. 43 in HALL, 1867. Dorsal, ventral, anterior, posterior, and lateral views. Costal formula:  $\frac{5}{4}$ ;  $\frac{1-1}{1-1}$ ;  $\frac{7}{8}$  and  $\frac{8}{9}$ .
- Figs 16-20 — *Sapphicorhynchus sappho* (HALL, 1860). Paralectotype D, AMNH34625 = pl. 54, fig. 37 in HALL, 1867. Dorsal, ventral, anterior, posterior, and lateral views. Costal formula:  $\frac{6}{5}$ ;  $\frac{1-0}{1-0}$ ;  $\frac{7}{8}$ .
- Figs 21-25 — *Sapphicorhynchus sappho* (HALL, 1860). Paralectotype C, AMNH34624 = pl. 54, fig. 36 in HALL, 1867. Dorsal, ventral, anterior, posterior, and lateral views. Costal formula:  $\frac{5}{4}$ ;  $\frac{1-1}{1-1}$ ;  $\frac{6}{7}$ .
- Figs 26-30 — *Sapphicorhynchus sappho* (HALL, 1860). Paralectotype B, AMNH34623 = pl. 54, figs 34, 35 in HALL, 1867. Dorsal, ventral, anterior, posterior, and lateral views. Costal formula:  $\frac{6}{5}$ ; 0;  $\frac{7}{8}$  and  $\frac{8}{9}$ .
- Figs 31-33 — *Sapphicorhynchus sappho* (HALL, 1860). Paralectotype G, AMNH34628. Dorsal, ventral, anterior, posterior, and lateral views. Costal formula:  $\frac{5}{4}$ ;  $\frac{1-1}{1-1}$ ;  $\frac{7}{8}$ .
- Fig. 34 — *Rhynchonella* (*Stenocisma*) *sappho* var. AMNH34634 and 34631 = pl. 55, figs 51, 48 in HALL, 1867.






PLATE 1



